

The maximum likelihood degree of a very affine variety

June Huh

ABSTRACT

We show that the maximum likelihood degree of a smooth very affine variety is equal to the signed topological Euler characteristic. This generalizes Orlik and Terao's solution to Varchenko's conjecture on complements of hyperplane arrangements to smooth very affine varieties. For very affine varieties satisfying a genericity condition at infinity, the result is further strengthened to relate the variety of critical points to the Chern-Schwartz-MacPherson class. The strengthened version recovers the geometric deletion-restriction formula of Denham, Garroubian, and Schulze for arrangement complements, and generalizes Kouchnirenko's theorem on the Newton polytope for nondegenerate hypersurfaces.

1. Introduction

Maximum likelihood estimation in statistics leads to the problem of finding critical points of a product of powers of polynomials on an algebraic variety [PS05, Section 3.3]. When the polynomials and the variety are linear and defined over the real numbers, the number of critical points is the number of bounded regions in the corresponding arrangement of hyperplanes.

Studying Bethe vectors in statistical mechanics, Varchenko conjectured a combinatorial formula for the number of critical points for complex hyperplane arrangements [Var95]. Let U be the complement of n hyperplanes in \mathbb{C}^r defined by the linear functions f_1, \dots, f_n . The master function $\varphi_{\mathbf{u}} = \prod_{i=1}^n f_i^{u_i}$, where the exponents u_i are integral parameters, is a holomorphic function on U . We assume that the affine hyperplane arrangement is *essential*, meaning that the lowest dimensional intersections of the hyperplanes are isolated points.

VARCHENKO'S CONJECTURE. *If the hyperplane arrangement is essential and the exponents u_i are sufficiently general, then the following holds:*

- (i) $\varphi_{\mathbf{u}}$ has only finitely many critical points in U .
- (ii) All critical points of $\varphi_{\mathbf{u}}$ are nondegenerate.
- (iii) The number of critical points is equal to the signed Euler characteristic $(-1)^r \chi(U)$.

The conjecture is proved by Varchenko when the hyperplanes are defined over the real numbers [Var95], and by Orlik and Terao in general [OT95]. Subsequent works of Silvotti and Damon extended this result to some nonlinear arrangements [Dam99, Dam00, Sil96]. The assumption made on the arrangement is certainly necessary, for there are arrangements violating the inequality $(-1)^r \chi(U) \geq 0$.

The principal aim of this paper is to generalize the theorem of Orlik and Terao. The generalization is pursued in two directions. In Theorem 1, we obtain the same conclusion for a wider class of affine varieties than complements of essential arrangements, and in Theorem 2, we recover the whole characteristic class from the critical points instead of the topological Euler characteristic. A connection to Kouchnirenko's theorem on the relation between the Newton polytope and the Euler characteristic is pointed out in Section 4.

The above extensions are motivated by the problem of maximum likelihood estimation in algebraic statistics. Recall that an irreducible algebraic variety is called *very affine* if it is isomorphic to a closed subvariety of an algebraic torus. Very affine varieties recently received a considerable amount of attention due to their central role in tropical geometry [EKL06, Spe05, Tev07]. The complement of an affine hyperplane arrangement is affine, and it is very affine if and only if the hyperplane arrangement is essential. Any complement of an affine hyperplane arrangement is of the form $U \times \mathbb{C}^k$, where U is the complement of an essential arrangement.

In view of the maximum likelihood estimation, very affine varieties are natural class of objects generalizing complements of hyperplane arrangements. Consider the projective space with the homogeneous coordinates p_1, \dots, p_n , where the coordinate p_i represents the probability of the i -th event. An implicit statistical model is a closed subvariety $V \subseteq \mathbb{P}^{n-1}$. The data comes in the form of nonnegative integers u_1, \dots, u_n , where u_i is the number of times the i -th event was observed.

In order to find the values of p_i on V which best explains the given data u_i , one finds critical points of the likelihood function

$$L(p_1, \dots, p_n) = p_1^{u_1} \cdots p_n^{u_n} / (p_1 + \cdots + p_n)^{u_1 + \cdots + u_n}.$$

Statistical computations are typically done in the affine chart defined by the nonvanishing of $p_1 + \cdots + p_n$, where the sum can be set equal to 1 and the denominator of L can be ignored. The maximum likelihood degree of the model is defined to be the number of complex critical points of the restriction of L to the projective variety V , where we only count critical points that are not poles or zeros of L , and u_1, \dots, u_n are assumed to be sufficiently general [HKS05]. In other words, the maximum likelihood degree is the number of critical points of the likelihood function on the very affine variety

$$U := \{x \in V \mid p_1 \cdots p_n(p_1 + \cdots + p_n) \neq 0\}.$$

1.1 Varchenko's conjecture for very affine varieties

We extend the theorem of Orlik and Terao to smooth very affine varieties. Let U be a smooth very affine variety of dimension r . Choose a closed embedding

$$f : U \longrightarrow (\mathbb{C}^*)^n, \quad f = (f_1, \dots, f_n).$$

The master function $\varphi_{\mathbf{u}} = \prod_{i=1}^n f_i^{u_i}$, where the exponents u_i are integral parameters, is a holomorphic function on U . The *maximum likelihood degree* of U is defined to be the number of critical points of the master function with sufficiently general exponents u_i .

THEOREM 1. *If the exponents u_i are sufficiently general, then the following holds:*

- (i) $\varphi_{\mathbf{u}}$ has only finitely many critical points in U .
- (ii) All critical points of $\varphi_{\mathbf{u}}$ are nondegenerate.
- (iii) The number of critical points is equal to the signed Euler characteristic $(-1)^r \chi(U)$.

More precisely, there is a nonzero polynomial F such that the assertions are valid for u_1, \dots, u_n with $F(u_1, \dots, u_n) \neq 0$.

Theorem 1 shows that, for example, the conclusions of [CHKS06, Theorem 20] and [Dam99, Corollary 6] hold for smooth very affine varieties without further assumptions. This has a few immediate corollaries that might be of interest in algebraic geometry and algebraic statistics. First, the maximum likelihood degree does not depend on the embedding of U into an algebraic torus. Second, the maximum likelihood degree satisfies the deletion-restriction formula as in the case of a linear model. Third, the sign of the Euler characteristic of a smooth very affine variety depends only on the parity of its dimension.

1.2 A geometric formula for the CSM class

The theorem of Orlik and Terao can be further generalized for very affine varieties which admit a good tropical compactification in the sense of Tevelev [Tev07]. See Definition 3.6 for *schön* very affine varieties. For example, the complement of an essential hyperplane arrangement is schön, and the hypersurface defined by a sufficiently general Laurent polynomial (with respect to its Newton polytope) is schön. The open subset of the grassmannian $\mathrm{Gr}(2, n)$ given by nonvanishing of all Plücker coordinates is another schön very affine variety of interest in algebraic statistics [SS04].

The generalization is formulated in terms of the variety of critical points of U , the totality of critical points of all possible (multivalued) master functions for U . More precisely, given a compactification \overline{U} of U , the variety of critical points $\mathfrak{X}(U)$ is defined to be the closure

$$\mathfrak{X}(U) = \overline{\mathfrak{X}^\circ(U)} \subseteq \overline{U} \times \mathbb{P}^{n-1}, \quad \mathfrak{X}^\circ(U) = \left\{ \sum_{i=1}^n u_i \cdot \mathrm{dlog}(f_i)(x) = 0 \right\} \subseteq U \times \mathbb{P}^{n-1},$$

where \mathbb{P}^{n-1} is the projective space with the homogeneous coordinates u_1, \dots, u_n . The variety of critical points has been previously studied in the context of hyperplane arrangements [CDFV11, DGS12]. See Section 2 for a detailed construction in the general setting.

We relate the variety of critical points to the Chern-Schwartz-MacPherson class of [Mac74]. Let \mathbb{T}_U be the intrinsic torus of U , an algebraic torus containing U whose character lattice is the group of nonvanishing regular functions on U modulo nonzero constants. We compactify the intrinsic torus by the projective space \mathbb{P}^n , where n is the dimension of \mathbb{T}_U .

THEOREM 2. *Suppose U is an r -dimensional very affine variety not isomorphic to a torus. If U is schön, then*

$$[\mathfrak{X}(U)] = \sum_{i=0}^r v_i [\mathbb{P}^{r-i} \times \mathbb{P}^{n-1-r+i}] \in A_*(\mathbb{P}^n \times \mathbb{P}^{n-1}),$$

where

$$c_{SM}(\mathbf{1}_U) = \sum_{i=0}^r (-1)^i v_i [\mathbb{P}^{r-i}] \in A_*(\mathbb{P}^n).$$

Theorem 1 is recovered by considering the number of points in a general fiber of the second projection from $\mathfrak{X}(U)$, which is the maximum likelihood degree

$$v_r = (-1)^r \int c_{SM}(\mathbf{1}_U) = (-1)^r \chi(U).$$

When U is the complement of an essential hyperplane arrangement \mathcal{A} and \mathbb{P}^n is the usual com-

pactification of \mathbb{T}_U defined by the ratios of homogeneous coordinates $z_1/z_0, \dots, z_n/z_0$, Theorem 2 specializes to the geometric formula for the characteristic polynomial of Denham, Garroubian, and Schulze [DGS12, Theorem 1.1]:

$$\chi_{\mathcal{A}}(q+1) = \sum_{i=0}^r (-1)^i v_i q^{r-i}.$$

The formula is used in [Huh12b] to verify the conjecture of Dawson on the logarithmic concavity of the h -vector of a matroid complex, for matroids representable over a field of characteristic zero [Daw84]. Other implications of the geometric formula are collected in Remark 3.12.

1.3 A generalization of Kouchnirenko's theorem

A schön hypersurface in an algebraic torus is defined by a Laurent polynomial which is nondegenerate in the sense of Kouchnirenko [Kou76]. We generalize Kouchnirenko's theorem equating the Euler characteristic with the signed volume of the Newton polytope, in the setting of Theorem 2. We hope that the approach of the present paper clarifies an analogy noted in [Var95, Remarks (e)], where Varchenko asks for a connection between Kouchnirenko's theorem and the conjecture stated in the introduction.

Let g be a Laurent polynomial in n variables with the Newton polytope Δ_g , and denote the corresponding hypersurface by

$$U = \{g = 0\} \subseteq (\mathbb{C}^*)^n.$$

Fix the open embedding $(\mathbb{C}^*)^n \subset \mathbb{P}^n$ defined by the ratios homogeneous coordinates $z_1/z_0, \dots, z_n/z_0$.

We follow the convention of [CLO98, Chapter 7] and write MV_n for the n -dimensional mixed volume. For example, the n -dimensional standard simplex Δ in \mathbb{R}^n has the unit volume 1.

THEOREM 3. *Let g be a nonzero Laurent polynomial in $n = r + 1$ variables with*

$$c_{SM}(\mathbf{1}_U) = \sum_{i=0}^r (-1)^i v_i [\mathbb{P}^{r-i}] \in A_*(\mathbb{P}^n).$$

If g is nondegenerate, then

$$v_i = MV_n(\underbrace{\Delta, \dots, \Delta}_{r-i}, \underbrace{\Delta_g, \dots, \Delta_g}_{i+1}), \quad i = 0, \dots, r.$$

In particular, the maximum likelihood degree of U is equal to the normalized volume

$$v_r = (-1)^r \int c_{SM}(\mathbf{1}_U) = \text{Volume}(\Delta_g).$$

Theorem 3 has applications not covered by Kouchnirenko's theorem. In particular, we get an explicit formula for the degree of the gradient map of a homogeneous polynomial in terms of the Newton polytope. See Corollary 4.4. This shows that many delicate examples discovered in classical projective geometry have a rather simple combinatorial origin.

As an example, we find irreducible homaloidal projective hypersurfaces of the given degree $d \geq 3$ and the ambient dimension $n \geq 3$, improving previous constructions [CRS08, FM12]. See Example 4.7.

1.4 Organization

We provide a brief overview of the paper.

Section 2 is devoted to the proof of Theorem 1. Along the way we construct the variety of critical points and describe its basic properties.

Section 3 introduces the deletion-restriction for very affine varieties, extending that for hyperplane arrangements. A brief introduction to the Chern-Schwartz-MacPherson class is given, and Theorem 2 is proved.

Section 4 focuses on the maximum likelihood degree of nondegenerate hypersurfaces in algebraic tori. Applications of Theorem 3 to the geometry of projective hypersurfaces are given.

ACKNOWLEDGEMENTS

The author is grateful to Bernd Sturmfels for asking the right questions which initiated this paper. He thanks Mircea Musta a for useful discussions throughout the preparation of the manuscript. He thanks Nero Budur, Eric Katz, and Sam Payne for helpful comments.

2. Proof of Theorem 1

2.1 The Gauss map of very affine varieties

An important role will be played by the Gauss map of a very affine variety in its intrinsic torus. Let U be a smooth very affine variety of dimension r . Choose a closed embedding

$$f : U \longrightarrow (\mathbb{C}^*)^n, \quad f = (f_1, \dots, f_n).$$

By a theorem of Samuel [Sam66], the group of invertible regular functions $M_U := \mathbb{C}[U]^*/\mathbb{C}^*$ is a finitely generated free abelian group. Therefore one may choose f_i to form a basis of M_U . In this case, f is a closed embedding of U into the intrinsic torus \mathbb{T}_U with the character lattice M_U

$$f : U \longrightarrow \mathbb{T}_U.$$

Any morphism from U to an algebraic torus \mathbb{T} is a composition of f with a homomorphism $\mathbb{T}_U \rightarrow \mathbb{T}$. The Gauss map of U is defined by the pushforward of f followed by the left-translation to the identity

$$T_x U \longrightarrow T_{f(x)} \mathbb{T}_U \longrightarrow T_1 \mathbb{T}_U, \quad x \in U.$$

In coordinates, the first map is represented by the jacobian matrix

$$\left(\frac{\partial f_i}{\partial x_j} \right), \quad 1 \leq i \leq n, \quad 1 \leq j \leq r,$$

and the second map is represented by the diagonal matrix with the diagonal entries $1/f_i(x)$. The composition of the two is the logarithmic jacobian matrix

$$\left(\frac{\partial \log f_i}{\partial x_j} \right), \quad 1 \leq i \leq n, \quad 1 \leq j \leq r.$$

This defines the Gauss map from U to the Grassmannian of $T_1 \mathbb{T}_U$

$$U \longrightarrow \mathrm{Gr}_r(T_1 \mathbb{T}_U), \quad x \longmapsto T_x U \subseteq T_1 \mathbb{T}_U.$$

Let Ω_U^1 be the sheaf of differential one-forms on U . Consider the complex vector space

$$W := M_U \otimes_{\mathbb{Z}} \mathbb{C}.$$

The dependence of W on U will often be omitted from the notation. We write W_U for (the sheaf sections of) the trivial vector bundle over U with the fiber W . There is a vector bundle homomorphism Φ defined by the evaluation of the logarithmic differential forms

$$\Phi : W_U \longrightarrow \Omega_U^1, \quad \left(\prod_{i=1}^n f_i^{u_i}, x \right) \longmapsto \sum_{i=1}^n u_i \cdot \mathrm{dlog}(f_i)(x).$$

At a point $x \in U$, the linear map $\Phi(x)$ between the fibers is dual to the injective linear map considered in Section 2.1:

$$T_x U \longrightarrow T_{f(x)} \mathbb{T}_U \longrightarrow T_1 \mathbb{T}_U.$$

Therefore Φ is surjective and $\ker \Phi$ is a vector bundle over U .

2.2 The variety of critical points

The inclusion of $\ker \Phi$ into W_U defines a closed immersion between the projective bundles

$$\mathfrak{X}^\circ(U) := \mathrm{Proj}(\mathrm{Sym}(\ker \Phi^\vee)) \longrightarrow \mathrm{Proj}(\mathrm{Sym}(W_U^\vee)) \simeq U \times \mathbb{P}(W).$$

Note that the following conditions are equivalent:

1. Φ is injective.
2. U is isomorphic to a torus.
3. $\mathfrak{X}^\circ(U)$ is empty.

If $\mathfrak{X}^\circ(U)$ is not empty, then $\mathfrak{X}^\circ(U)$ is a projective bundle over U of dimension equal to that of $\mathbb{P}(W)$, defined by the equation

$$\sum_{i=1}^n u_i \cdot \mathrm{dlog}(f_i)(x) = 0,$$

where u_1, \dots, u_n are the homogeneous coordinates of $\mathbb{P}(W)$. In short, $\mathfrak{X}^\circ(U)$ is the set of critical points of all possible (multivalued) master functions.

DEFINITION 2.1. Given a compactification V of U , the *variety of critical points* of U is defined to be the closure

$$\mathfrak{X}_V(U) := \overline{\mathfrak{X}^\circ(U)} \subseteq V \times \mathbb{P}(W).$$

We denote the variety of critical points by $\mathfrak{X}(U)$ when there is no danger of confusion.

The variety of critical points is irreducible by construction. When U is the complement of an essential arrangement of hyperplanes, $\mathfrak{X}(U)$ is the variety of critical points previously considered in the context of hyperplane arrangements [CDFV11, DGS12]. This variety has its origin in [OT95, Proposition 4.1].

We record here the following basic compatibility: If $V_1 \rightarrow V_2$ is a morphism between two compactifications of U which is the identity on U , then the class of $\mathfrak{X}_{V_1}(U)$ maps to the class of $\mathfrak{X}_{V_2}(U)$ under the induced map between the Chow groups

$$A_*(V_1 \times \mathbb{P}(W)) \longrightarrow A_*(V_2 \times \mathbb{P}(W)), \quad [\mathfrak{X}_{V_1}(U)] \longmapsto [\mathfrak{X}_{V_2}(U)].$$

Remark 2.2. We point out a technical difference between Definition 2.1 and the variety of critical points in the cited literature. This remark is intended for readers familiar with [CDFV11, DGS12], and is independent from the rest of the paper.

Let U be the complement of \mathcal{A} , an essential arrangement of n hyperplanes in \mathbb{C}^r . The variety of critical points in [DGS12] is defined when \mathcal{A} is a central arrangement, so we suppose that this is the case. In our notation, it is the quotient under the torus action

$$\tilde{\mathfrak{X}}(U) := \mathfrak{X}_V(U)/(\mathbb{C}^* \times \mathbf{1}) \subseteq (\mathbb{C}^r \times \mathbb{P}^{n-1})/(\mathbb{C}^* \times \mathbf{1}) = \mathbb{P}^{r-1} \times \mathbb{P}^{n-1},$$

where V is the partial compactification of U by the affine space \mathbb{C}^r . Since \mathcal{A} is central, by [OT95, Proposition 3.9],

$$\tilde{\mathfrak{X}}(U) \subseteq \{u_1 + \cdots + u_n = 0\} \simeq \mathbb{P}^{r-1} \times \mathbb{P}^{n-2} \subset \mathbb{P}^{r-1} \times \mathbb{P}^{n-1}.$$

The quotient variety $\tilde{\mathfrak{X}}(U)$ indeed is the variety of critical points (of a closely related arrangement) in our sense. Consider a decone $\tilde{\mathcal{A}}$ of \mathcal{A} , an affine arrangement obtained by declaring one of the hyperplanes in the projectivization of \mathcal{A} to be the hyperplane at infinity. The number of hyperplanes and the rank of the decone are one less than those of \mathcal{A} . More precisely, we have the relation between the characteristic polynomials

$$\chi_{\tilde{\mathcal{A}}}(q) = \chi_{\mathcal{A}}(q)/(q-1).$$

Let \tilde{U} be the complement of the decone in \mathbb{C}^{r-1} , and take the obvious compactification \mathbb{P}^{r-1} of \tilde{U} . Then the variety of critical points of \tilde{U} is the subvariety considered above:

$$\tilde{\mathfrak{X}}(U) \subseteq \mathbb{P}^{r-1} \times \mathbb{P}^{n-2}.$$

The reader is invited to compare the formula of Corollary 3.11 with its cohomology version [DGS12, Theorem 1.1] for essential central arrangements.

2.3 Nonvanishing at the boundary

Let H_0, \dots, H_n be the torus invariant hyperplanes in \mathbb{P}^n defined by the homogeneous coordinates z_0, \dots, z_n . Fix the open embedding

$$\iota : (\mathbb{C}^*)^n \longrightarrow \mathbb{P}^n,$$

defined by the ratios $z_1/z_0, \dots, z_n/z_0$. Let V be the closure of U in \mathbb{P}^n , and choose a simple normal crossing resolution of singularities

$$\begin{array}{ccc} \pi^{-1}(U) & \longrightarrow & \tilde{V} \\ \downarrow & & \downarrow \pi \\ U & \xrightarrow{\iota} & V \longrightarrow \mathbb{P}^n, \end{array}$$

where π is an isomorphism over U , \tilde{V} is smooth, and $\tilde{V} \setminus \pi^{-1}(U)$ is a simple normal crossing divisor with the irreducible components D_1, \dots, D_k . Our goal is to show that a sufficiently general differential form on \tilde{V} with logarithmic poles along D_j has a zero scheme which is a finite set of reduced points in U .

Each f_i defines a rational function on \tilde{V} which is regular on $\pi^{-1}(U)$. We have

$$\text{ord}_{D_j}(f_i) = \begin{cases} \text{positive} & \text{if } \pi(D_j) \not\subseteq H_0 \text{ and } \pi(D_j) \subseteq H_i, \\ \text{negative} & \text{if } \pi(D_j) \subseteq H_0 \text{ and } \pi(D_j) \not\subseteq H_i. \end{cases}$$

LEMMA 2.3. *For each j , there is an i such that $\text{ord}_{D_j}(f_i)$ is nonzero.*

Proof. Since D_j is irreducible, each $\pi(D_j)$ is contained in some H_i . The assertion follows from the following set-theoretic reasoning:

1. If $\pi(D_j) \not\subseteq H_0$, then $\pi(D_j) \subseteq H_i$ for some i because $\bigcup_{i=0}^n H_i = \mathbb{P}^n \setminus (\mathbb{C}^*)^n$.
2. If $\pi(D_j) \subseteq H_0$, then $\pi(D_j) \not\subseteq H_i$ for some i because $\bigcap_{i=0}^n H_i = \emptyset$.

□

An integral vector $\mathbf{u} = (u_1, \dots, u_n) \in \mathbb{Z}^n$ defines a rational function on \tilde{V}

$$\varphi_{\mathbf{u}} = \prod_{i=1}^n f_i^{u_i},$$

which is regular on $\pi^{-1}(U)$. Note that for each j ,

$$\text{ord}_{D_j}(\varphi_{\mathbf{u}}) = \sum_{i=1}^n u_i \cdot \text{ord}_{D_j}(f_i).$$

Combining this with Lemma 2.3, we have

LEMMA 2.4. *For a sufficiently general $\mathbf{u} \in \mathbb{Z}^n$, $\text{ord}_{D_j}(\varphi_{\mathbf{u}})$ is nonzero for all $j = 1, \dots, k$.*

More precisely, there are corank 1 subgroups A_1, \dots, A_k of \mathbb{Z}^n such that $\text{ord}_{D_j}(\varphi_{\mathbf{u}})$ is nonzero for $\mathbf{u} \in \mathbb{Z}^n \setminus \bigcup_{j=1}^k A_j$ and $j = 1, \dots, k$.

Consider the sheaf of logarithmic differential one-forms $\Omega_{\tilde{V}}^1(\log D)$, where $D = D_1 + \dots + D_k$. For the definition and the needed properties of the sheaf of logarithmic differential one-forms, we refer to [Del70, Sai80]. $\Omega_{\tilde{V}}^1(\log D)$ is a locally free sheaf of rank r , and the rational function $\varphi_{\mathbf{u}}$ defines a global section

$$\text{dlog}(\varphi_{\mathbf{u}}) = \sum_{i=1}^n u_i \cdot \text{dlog}(f_i) \in H^0(\tilde{V}, \Omega_{\tilde{V}}^1(\log D)).$$

LEMMA 2.5. *For a sufficiently general $\mathbf{u} \in \mathbb{Z}^n$, $\text{dlog}(\varphi_{\mathbf{u}})$ does not vanish on D .*

Proof. Given a point $x \in D$, let D_1, \dots, D_l be the irreducible components of D containing x , and let g_1, \dots, g_l be local defining equations on a small neighborhood G of x . Clearly, l is at least one. By replacing G to a smaller neighborhood if necessary, we may assume that $\Omega_{\tilde{V}}^1(\log D)$ trivializes over G , and

$$\varphi_{\mathbf{u}} = g_1^{o_1} \cdots g_l^{o_l} h, \quad o_j = \text{ord}_{D_j}(\varphi_{\mathbf{u}}),$$

for some nonvanishing holomorphic function h on G . Over the open set G , we have

$$\text{dlog}(\varphi_{\mathbf{u}}) = \left(\sum_{j=1}^l \text{ord}_{D_j}(\varphi_{\mathbf{u}}) \cdot \text{dlog}(g_j) \right) + \psi,$$

where ψ is a regular differential one-form. Since $\text{dlog}(g_j)$ form a part of a free basis of a trivialization of $\Omega_{\tilde{V}}^1(\log D)$ over G , it follows from Lemma 2.4 that $\text{dlog}(\varphi_{\mathbf{u}})$ does not vanish on D for a sufficiently general $\mathbf{u} \in \mathbb{Z}^n$. □

2.4 Proof of Theorem 1

Let $\mathfrak{X}(U) = \mathfrak{X}_{\tilde{V}}(U)$ be the variety of critical points of U in $\tilde{V} \times \mathbb{P}(W)$, where W is the complex vector space $M_U \otimes_{\mathbb{Z}} \mathbb{C}$ as before. Write $W_{\tilde{V}}$ for (the sheaf sections of) the trivial vector bundle over \tilde{V} with the fiber W , and consider the homomorphism Ψ defined by the evaluation of the

logarithmic differential forms

$$\Psi : W_{\tilde{V}} \longrightarrow \Omega_{\tilde{V}}^1(\log D), \quad (x, \mathbf{u}) \longmapsto \mathrm{dlog}(\varphi_{\mathbf{u}})(x) := \sum_{i=1}^n u_i \cdot \mathrm{dlog}(f_i)(x).$$

We do not attempt to give an individual meaning to the multivalued master function $\varphi_{\mathbf{u}}$ when $\mathbf{u} = (u_1, \dots, u_n)$ is not integral. Let $\mathrm{pr}_1, \mathrm{pr}_2$ be the two projections from $\tilde{V} \times \mathbb{P}(W)$, and define the incidence variety of the evaluation

$$\mathcal{J}(U) = \mathcal{J}_{\tilde{V}}(U) = \{(x, \mathbf{u}) \in \tilde{V} \times \mathbb{P}(W) \mid \mathrm{dlog}(\varphi_{\mathbf{u}})(x) = 0\}.$$

We drop the subscript from $\mathcal{J}_{\tilde{V}}(U)$ when there is no danger of confusion. By definition, $\mathfrak{X}(U)$ is the unique irreducible component of $\mathcal{J}(U)$ which dominates \tilde{V} under pr_1 .

Then, for a sufficiently general $\mathbf{u} \in \mathbb{Z}^n$,

1. $\mathrm{pr}_2^{-1}(\mathbf{u}) \cap \mathcal{J}(U)$ is contained in $\mathfrak{X}^\circ(U)$, by Lemma 2.5, and
2. $\mathrm{pr}_2^{-1}(\mathbf{u}) \cap \mathfrak{X}^\circ(U)$ is a finite set of reduced points, by the Bertini theorem applied to pr_2 on $\mathfrak{X}(U)$ [Jou83, Theoreme 6.10].

More precisely, there is a nonempty Zariski open subset $\mathcal{U} \subseteq W$ such that the two assertions are valid for any element in the infinite set $M_U \cap \mathcal{U}$.

It follows that the zero scheme of the section $\mathrm{dlog}(\varphi_{\mathbf{u}}) \in H^0(\tilde{V}, \Omega_{\tilde{V}}^1(\log D))$,

$$\mathrm{pr}_1(\mathrm{pr}_2^{-1}(\mathbf{u})) = \{x \in \tilde{V} \mid \mathrm{dlog}(\varphi_{\mathbf{u}})(x) = 0\},$$

is a finite set of reduced points in U . The smoothness of \tilde{V} implies that the section $\mathrm{dlog}(\varphi_{\mathbf{u}})$ is regular, and the above set represents the homology class $c_r(\Omega_{\tilde{V}}^1(\log D)) \cap [\tilde{V}]$ [Ful98, Example 3.2.16]. Since all the f_i are nonvanishing on U , we may identify the critical points of $\varphi_{\mathbf{u}}$ with the zero scheme of $\mathrm{dlog}(\varphi_{\mathbf{u}})$.

Therefore all the critical points of $\varphi_{\mathbf{u}}$ are nondegenerate, and the number of critical points is equal to the degree of the top Chern class of $\Omega_{\tilde{V}}^1(\log D)$. Finally, from the logarithmic Poincaré-Hopf theorem [Kaw78, Nor78, Sil96], we have

$$\int_{\tilde{V}} c_r(\Omega_{\tilde{V}}^1(\log D)) = (-1)^r \int_{\tilde{V}} c_r(\Omega_{\tilde{V}}^1(\log D)^\vee) = (-1)^r \chi(U).$$

□

3. Deletion-restriction for very affine varieties

In this section we formulate the deletion-restriction for the characteristic polynomial of a hyperplane arrangement in the very affine setting. The role of the characteristic polynomial will be played by a characteristic class for very affine varieties.

This point of view is particularly satisfactory for very affine varieties satisfying a genericity condition at infinity, and gives new insights on the positivity of the coefficients of the characteristic polynomial. For complements of hyperplane arrangements, we recover the geometric formula for the characteristic polynomial of Denham, Garrounian, and Schulze [DGS12, Theorem 1.1].

Let U be a very affine variety, and let U_0 be the hypersurface of U defined by the vanishing of a regular function. The complement $U_1 = U \setminus U_0$ is a very affine variety, being a principal affine open subset of a very affine variety.

DEFINITION 3.1. A *triple* of very affine varieties is a collection of the form (U, U_0, U_1) .

In the language of hyperplane arrangements, U corresponds to the arrangement obtained by deleting the distinguished hyperplane U_0 from the arrangement corresponding to U_1 . For this reason, we call U the deletion of U_1 , and U_0 the restriction of U_1 . As in the case of hyperplane arrangements, we have

$$\dim U_1 = \dim U \quad \text{and} \quad \dim U_0 = \dim U - 1.$$

By the set-theoretic additivity of the topological Euler characteristics of complex algebraic varieties [Ful93, Section 4.5], we have

$$\chi(U) = \chi(U_1) + \chi(U_0).$$

Therefore, by Theorem 1, the maximum likelihood degrees of a triple of smooth very affine varieties satisfy an additive formula. Write $ML(U)$ for the maximum likelihood degree of a very affine variety U , the number of critical points of a master function of U with sufficiently general exponents.

COROLLARY 3.2. *If (U, U_0, U_1) is a triple of smooth very affine varieties, then*

$$ML(U) = ML(U_1) - ML(U_0).$$

It would be interesting to obtain a direct justification of Corollary 3.2.

Remark 3.3. Let us call a very affine variety *primitive* if it does not have any deletion. For example, the complement of an essential hyperplane arrangement is primitive if and only if it is the complement of a Boolean arrangement. Note that this is the case exactly when the complement is isomorphic to its intrinsic torus. A distinguished feature of the deletion-restriction for very affine varieties, when compared to that for hyperplane arrangements, is that there are primitive very affine varieties not isomorphic to a torus. These very affine varieties are responsible for the noncombinatorial aspect of the extended theory.

When (U, U_0, U_1) is a triple of hyperplane arrangement complements, Corollary 3.2 is the deletion-restriction formula for the Möbius invariant $\chi_{\mathcal{A}}(1)$, where $\chi_{\mathcal{A}}(q)$ is the characteristic polynomial of an affine hyperplane arrangement \mathcal{A} . The full deletion-restriction formula between the characteristic polynomials can be formulated in terms of the Chern-Schwartz-MacPherson class [Mac74]. Below we give a brief description of the CSM class; Aluffi provides a gentle introduction in [Alu05].

Recall that the group of constructible functions $C(X)$ of an algebraic variety X is generated by functions of the form $\mathbf{1}_Z$, where Z is a subvariety of X . If $f : X \rightarrow Y$ is a morphism between complex algebraic varieties, then the pushforward of constructible functions is defined by the homomorphism

$$f_* : C(X) \longrightarrow C(Y), \quad \mathbf{1}_Z \longmapsto \left(p \longmapsto \chi(f^{-1}(p) \cap Z) \right).$$

If X is a compact complex manifold, then the characteristic class of X is the Chern class of the tangent bundle $c(TX) \cap [X] \in A_*(X)$, where $A_*(X)$ is the Chow homology group of X [Ful98]. A generalization is provided by the Chern-Schwartz-MacPherson class, whose existence was once a conjecture of Deligne and Grothendieck. For a construction with emphasis on smooth and possibly noncompact varieties, see [Alu06b].

Let C be the functor of constructible functions from the category of complex algebraic varieties (with proper morphisms) to the category of abelian groups.

DEFINITION 3.4. The *CSM class* is the unique natural transformation

$$c_{SM} : C \longrightarrow A_*$$

such that $c_{SM}(\mathbf{1}_X) = c(TX) \cap [X]$ when X is smooth and complete.

The uniqueness follows from the naturality, the resolution of singularities, and the normalization for smooth and complete varieties. The CSM class satisfies the inclusion-exclusion

$$c_{SM}(\mathbf{1}_{U \cup U'}) = c_{SM}(\mathbf{1}_U) + c_{SM}(\mathbf{1}_{U'}) - c_{SM}(\mathbf{1}_{U \cap U'}),$$

and captures the Euler characteristic as its degree

$$\chi(U) = \int c_{SM}(\mathbf{1}_U).$$

When U is the complement of an arrangement of hyperplanes \mathcal{A} in \mathbb{C}^r , the CSM class of U is the characteristic polynomial $\chi_{\mathcal{A}}(q)$. For the definition of the characteristic polynomial of an affine arrangement, see [OT92, Definition 2.52].

THEOREM 3.5. *Let \mathbb{P}^r be the compactification of U defined by the hyperplane at infinity $\mathbb{P}^r \setminus \mathbb{C}^r$. Then*

$$c_{SM}(\mathbf{1}_U) = \sum_{i=0}^r (-1)^i v_i [\mathbb{P}^{r-i}] \in A_*(\mathbb{P}^r),$$

where

$$\chi_{\mathcal{A}}(q+1) = \sum_{i=0}^r (-1)^i v_i q^{r-i}.$$

This is because the recursive formula for a triple of arrangement complements

$$c_{SM}(\mathbf{1}_{U_1}) = c_{SM}(\mathbf{1}_U - \mathbf{1}_{U_0}) = c_{SM}(\mathbf{1}_U) - c_{SM}(\mathbf{1}_{U_0}),$$

agrees with the usual deletion-restriction formula [OT92, Theorem 2.56]

$$\chi_{\mathcal{A}_1}(q+1) = \chi_{\mathcal{A}}(q+1) - \chi_{\mathcal{A}_0}(q+1).$$

The induction is on the dimension and on the number of hyperplanes. The case of no hyperplanes is a direct computation of $c_{SM}(\mathbf{1}_{\mathbb{C}^r})$ by the inclusion-exclusion, and the case of dimension one is a special case of the equality

$$\chi(U) = \int c_{SM}(\mathbf{1}_U) = \chi_{\mathcal{A}}(1).$$

See [Alu12, Theorem 1.2] and also [Huh12a, Remark 26].

Our goal is to relate the variety of critical points to the CSM class. If we restrict our attention to the degree of the CSM class, then the relation recovers the conclusion stated in Varchenko's conjecture for the Euler characteristic. We prove this for a class of very affine varieties satisfying a genericity condition at infinity.

The genericity condition is commonly expressed using the language of tropical compactifications. If U is a subvariety of an algebraic torus \mathbb{T} , then we consider compactifications \overline{U} defined as the closures of U in various (not necessarily complete) toric varieties X of \mathbb{T} . We say that the compactification \overline{U} is *tropical* if the multiplication map

$$m : \mathbb{T} \times \overline{U} \longrightarrow X, \quad (t, x) \longmapsto tx$$

is flat and surjective. Tropical compactifications exist, and they are obtained from toric varieties X defined by sufficiently fine fan structures on the tropicalization of U [Tev07, Section 2].

DEFINITION 3.6. We say that U is *schön* if the multiplication is smooth for some tropical compactification of U .

Equivalently, U is schön if the multiplication is smooth for every tropical compactification of U [Tev07, Theorem 1.4].

Remark 3.7. There are two classes of schön very affine varieties of particular interest. The first is the class of complements of essential hyperplane arrangements, and the second is the class of nondegenerate hypersurfaces [Tev07]. What we need from the schön hypothesis is the existence of a simple normal crossings compactification which admits sufficiently many logarithmic differential one-forms. For arrangement complements, such a compactification is provided by the wonderful compactification of De Concini and Procesi [DP95]. For nondegenerate hypersurfaces, and more generally for nondegenerate complete intersections, the needed compactification has been constructed by Khovanskii [Hov77].

Let $U \subseteq \mathbb{T}_U$ be a very affine variety of dimension r , where \mathbb{T}_U is the intrinsic torus of Section 2.1. Let V be the closure of U in \mathbb{P}^n , where \mathbb{P}^n is a fixed toric compactification of \mathbb{T}_U . We follow Section 2.2 and define the variety of critical points

$$\mathfrak{X}(U) \subseteq V \times \mathbb{P}(W) \subseteq \mathbb{P}^n \times \mathbb{P}^{n-1}, \quad W = M_U \otimes_{\mathbb{Z}} \mathbb{C}.$$

THEOREM 3.8. Suppose U is schön and not isomorphic to a torus. Then

$$[\mathfrak{X}(U)] = \sum_{i=0}^r v_i [\mathbb{P}^{r-i} \times \mathbb{P}^{n-1-r+i}] \in A_*(\mathbb{P}^n \times \mathbb{P}^{n-1}),$$

where

$$c_{SM}(\mathbf{1}_U) = \sum_{i=0}^r (-1)^i v_i [\mathbb{P}^{r-i}] \in A_*(\mathbb{P}^n).$$

Proof. We prove a slightly more general statement. Let V be a compactification of U obtained by taking the closure in a toric variety X of \mathbb{T}_U . We will prove the equality

$$c_{SM}(\mathbf{1}_U) = \sum_{i=0}^r (-1)^i \text{pr}_{1*} [\text{pr}_2^{-1}(\mathbb{P}^{r-i}) \cap \mathfrak{X}(U)] \in A_*(V),$$

where pr_1, pr_2 are the two projections from $V \times \mathbb{P}(W)$, and \mathbb{P}^{r-i} is a sufficiently general linear subspace of $\mathbb{P}(W)$ of the indicated dimension. The projection formula shows that this implies the stated version when $X = \mathbb{P}^n$.

If U is a schön very affine variety, then there is a tropical compactification of U which has a simple normal crossings boundary divisor. More precisely, there is a smooth toric variety \tilde{X} of \mathbb{T}_U , obtained by taking a sufficiently fine fan structure on the tropicalization of U , such that the closure of U in \tilde{X} is a smooth and complete variety \tilde{V} with the simple normal crossings divisor $\tilde{V} \setminus U$ [Hac08, Proof of Theorem 2.5].

By taking a further subdivision of the fan of \tilde{X} if necessary, we may assume that there is a toric morphism $\tilde{X} \rightarrow X$ preserving \mathbb{T}_U . By the functoriality of the CSM class, we have

$$A_*(\tilde{V}) \longrightarrow A_*(V), \quad c_{SM}(\mathbf{1}_U) \longmapsto c_{SM}(\mathbf{1}_U).$$

Note also that

$$A_*(\tilde{V} \times \mathbb{P}(W)) \longrightarrow A_*(V \times \mathbb{P}(W)), \quad [\mathfrak{X}_{\tilde{V}}(U)] \longmapsto [\mathfrak{X}_V(U)].$$

By the projection formula, the problem is reduced to the case $\tilde{X} = X$.

In this case, we have the exact sequence induced by the restriction $\mathcal{O}_X|_V \rightarrow \mathcal{O}_V$

$$0 \longrightarrow N_{V/X}^\vee \longrightarrow \Omega_X^1(\log X \setminus \mathbb{T}_U)|_V \longrightarrow \Omega_V^1(\log V \setminus U) \longrightarrow 0.$$

Note that the middle term is isomorphic to the trivial vector bundle W_V over V with the fiber W [Ful93, Section 4.3]. Under this identification, the restriction of the differential one-forms is the evaluation map of Section 2.6:

$$0 \longrightarrow \ker \Psi \longrightarrow W_V \xrightarrow{\Psi} \Omega_V^1(\log V \setminus U).$$

It follows that the latter sequence is exact, and the projectivization of the kernel

$$\mathcal{J}(U) = \{(x, \mathbf{u}) \in V \times \mathbb{P}(W) \mid \mathrm{dlog}(\varphi_{\mathbf{u}})(x) = 0\}$$

coincides with the variety of critical points $\mathfrak{X}(U)$. Since the pullback of $\mathcal{O}_{\mathbb{P}(W)}(1)$ to $\mathfrak{X}(U)$ is the canonical line bundle $\mathcal{O}_{\mathbb{P}(N_{V/X}^\vee)}(1)$, we have

$$\begin{aligned} \sum_{i=0}^r \mathrm{pr}_{1*}[\mathrm{pr}_2^{-1}(\mathbb{P}^{r-i}) \cap \mathfrak{X}(U)] &= \mathrm{pr}_{1*} \left(\sum_{i=0}^r c_1(\mathrm{pr}_2^* \mathcal{O}_{\mathbb{P}(W)}(1))^{n-1-r+i} \cap [\mathfrak{X}(U)] \right) \\ &= s(N_{V/X}^\vee) \cap [V] = c(\Omega_V^1(\log V \setminus U)) \cap [V]. \end{aligned}$$

Here n is the dimension of W , and the last equality is the Whitney sum formula. Now the assertion follows from the fact that the CSM class of a smooth variety is the Chern class of the logarithmic tangent bundle:

$$c_{SM}(\mathbf{1}_U) = c(\Omega_V^1(\log V \setminus U)^\vee) \cap [V].$$

This follows from a construction of the CSM class, which is most natural from the point of view of this paper [Alu06b, Section 4]. For precursors, see [Alu99, Theorem 1] and also [GP02, Proposition 15.3]. \square

Remark 3.9. In a refined form, the above proof shows that the equality

$$c_{SM}(\mathbf{1}_U) = \sum_{i=0}^r (-1)^i \mathrm{pr}_{1*}[\mathrm{pr}_2^{-1}(\mathbb{P}^{r-i}) \cap \mathfrak{X}(U)]$$

holds in the \mathbb{T} -equivariant proChow group $\hat{A}_*^{\mathbb{T}}(\mathbb{T}_U)$ of [Alu06a, Alu06b]. This removes the dependence on the compactification from Theorem 2. It should not be expected that the equality holds in the ordinary proChow group of \mathbb{T}_U .

Remark 3.10. Theorem 3.8 may fail to hold for a smooth very affine variety. As an example, consider a smooth hypersurface U in $(\mathbb{C}^*)^3$ whose closure in \mathbb{P}^3 has a node at a torus orbit of codimension one. A direct computation on a simple normal crossings compactification of U shows that the incidence variety $\mathcal{J}(U)$ has a two-dimensional component other than $\mathfrak{X}(U)$, and hence the classes of $\mathcal{J}(U)$ and $\mathfrak{X}(U)$ are different in $A_2(\mathbb{P}^3 \times \mathbb{P}^2)$.

For complements of hyperplane arrangements, Theorem 3.8 gives a geometric formula for the characteristic polynomial [DGS12, Theorem 1.1]. Let U be the complement of an arrangement of n distinct hyperplanes

$$\mathcal{A} = \{f_1 = 0\} \cup \cdots \cup \{f_n = 0\} \subset \mathbb{C}^r.$$

Then U is a very affine variety if and only if \mathcal{A} is an essential arrangement, meaning that the lowest dimensional intersections of the hyperplanes of \mathcal{A} are isolated points. Indeed, taking one of the isolated points as the origin of \mathbb{C}^r and choosing r linearly independent hyperplanes intersecting at the point reveals U as a principal affine open subset of $(\mathbb{C}^*)^r$. For the converse, one writes U as a product $U' \times \mathbb{C}^k$, where U' is the complement of an essential arrangement, and note that the affine line \mathbb{C} does not admit a closed embedding into an algebraic torus.

Suppose from now on that \mathcal{A} is an essential arrangement. \mathcal{A} is a Boolean arrangement if and only if U is isomorphic to an algebraic torus. The equations of the hyperplanes define a closed embedding

$$f : U \longrightarrow (\mathbb{C}^*)^n \simeq \mathbb{T}_U, \quad f = (f_1, \dots, f_n).$$

The indicated isomorphism follows from the linear independence of the f_i in M_U . We fix the open embedding $(\mathbb{C}^*)^n \subset \mathbb{P}^n$ defined by the ratios of homogeneous coordinates $z_1/z_0, \dots, z_n/z_0$. Then the closure of U in \mathbb{P}^n is a linear subspace $\mathbb{P}^r \subset \mathbb{P}^n$. In this setting, a combination of Theorem 3.5 and Theorem 3.8 gives the following statement.

COROLLARY 3.11. *Suppose \mathcal{A} is not a Boolean arrangement. Then*

$$[\mathfrak{X}(U)] = \sum_{i=0}^r v_i [\mathbb{P}^{r-i} \times \mathbb{P}^{n-1-r+i}] \in A_*(\mathbb{P}^r \times \mathbb{P}^{n-1}),$$

where

$$\chi_{\mathcal{A}}(q+1) = \sum_{i=0}^r (-1)^i v_i q^{r-i}.$$

Remark 3.12. A sequence e_0, \dots, e_r of integers is said to be *log-concave* if $e_{i-1}e_{i+1} \leq e_i^2$ for all i , and is said to have *no internal zeros* if the indices of the nonzero elements are consecutive integers. Write a homology class $\xi \in A_k(\mathbb{P}^n \times \mathbb{P}^m)$ as the linear combination

$$\xi = \sum_i e_i [\mathbb{P}^{k-i} \times \mathbb{P}^i].$$

It can be shown that some multiple of ξ is the fundamental class of an irreducible subvariety if and only if the e_i form a log-concave sequence of nonnegative integers with no internal zeros [Huh12a, Theorem 21].

Therefore, by Theorem 3.8, the v_i of a schön very affine variety form a log-concave sequence of nonnegative integers with no internal zeros. In particular, the coefficients of $\chi_{\mathcal{A}}(q+1)$ form a sequence with the three properties. This strengthens the previous result that the coefficients of $\chi_{\mathcal{A}}(q)$ form a log-concave sequence [Huh12a, Theorem 3], and answers several questions on sequences associated to a matroid, for matroids representable over a field of characteristic zero:

- (i) Read's conjecture predicts that the coefficient of the chromatic polynomial of a graph form a unimodal sequence [Rea68]. This follows from the log-concavity of $\chi_{\mathcal{A}}(q)$ when \mathcal{A} is the graphic arrangement of a given graph [Huh12a].
- (ii) Hoggar's conjecture predicts that the coefficient of the chromatic polynomial of a graph form a strictly log-concave sequence [Hog74]. This follows from the log-concavity of $\chi_{\mathcal{A}}(q+1)$ when \mathcal{A} is the graphic arrangement of a given graph [Huh12b].
- (iii) Welsh's conjecture predicts that the f -vector of a matroid complex form a unimodal sequence [Wel69]. This follows from the log-concavity of $\chi_{\mathcal{A}}(q)$ when \mathcal{A} is an arrangement corresponding to the cofree extension of a given matroid [Len12].

- (iv) Dawson's conjecture predicts that the h -vector of a matroid complex form a log-concave sequence [Daw84]. This follows from the log-concavity of $\chi_{\mathcal{A}}(q+1)$ when \mathcal{A} is an arrangement corresponding to the cofree extension of a given matroid [Huh12b].

For details on the derivation of the above variations, see [Huh12b].

Remark 3.13. The characteristic class approach to Varchenko's conjecture and the generalized deletion-restriction has been pioneered by Damon for nonlinear arrangements on smooth complete intersections [Dam99, Dam00]. In fact, it can be shown that Damon's higher multiplicities are the degrees of the CSM class of the arrangement complement. See [Huh12a] for the connection between the two, for nonlinear arrangements on a projective space.

The CSM point of view successfully deals with several problems considered by Damon in [Dam99, Dam00]. In particular, an affirmative answer to the conjecture in [Dam00, Remark 2.6] follows from the product formula

$$c_{SM}(\mathbf{1}_{U_1} \otimes \mathbf{1}_{U_2}) = c_{SM}(\mathbf{1}_{U_1}) \otimes c_{SM}(\mathbf{1}_{U_2}) \in A_*(\overline{U_1} \times \overline{U_2}).$$

The above is a refinement of the product formula of Kwieciński [Kwi92], and can be viewed as a generalization of the product formula for the Euler characteristic

$$\chi(U_1 \times U_2) = \chi(U_1)\chi(U_2).$$

See [Alu06a, Théorème 4.1].

4. Nondegenerate hypersurfaces

A schön hypersurface in an algebraic torus is defined by a nondegenerate Laurent polynomial. A sufficiently general Laurent polynomial with the given Newton polytope is nondegenerate, and the corresponding hypersurface admits a tropical compactification which has a simple normal crossings boundary divisor [Hov77].

We show that the variety of critical points of a schön hypersurface is controlled by the Newton polytope of the defining polynomial. This gives a formula for the CSM class in terms of the Newton polytope, which specializes to Kouchnirenko's theorem equating the Euler characteristic with the signed volume of the Newton polytope [Kou76, Théorème IV].

Let g be a nonzero Laurent polynomial in n variables

$$g = \sum_{\mathbf{u}} c_{\mathbf{u}} \mathbf{x}^{\mathbf{u}} \in \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}].$$

We are interested in the CSM class of the very affine variety

$$U = \{g = 0\} \subseteq (\mathbb{C}^*)^n.$$

The *Newton polytope* of g , denoted Δ_g , is the convex hull of exponents $\mathbf{u} \in \mathbb{Z}^n$ with the nonzero coefficient $c_{\mathbf{u}}$. Write g_{σ} for the Laurent polynomial made up of those terms of g which lie in a face σ of the Newton polytope. We say that g is *nondegenerate* if dg_{σ} is nonvanishing on $\{g_{\sigma} = 0\}$ for every face of its Newton polytope.

We follow the convention of [CLO98, Chapter 7] and write MV_n for the n -dimensional mixed volume. For example, the n -dimensional standard simplex Δ in \mathbb{R}^n has the normalized volume 1.

In view of later applications to projective hypersurfaces, we state our result for a fixed compactification $(\mathbb{C}^*)^n \subset \mathbb{P}^n$, where the open embedding is defined by the ratios of homogeneous

coordinates $z_1/z_0, \dots, z_n/z_0$. The formulation for other toric compactifications, and extension to complete intersections, are left to the interested reader.

THEOREM 4.1. *Let g be a nonzero Laurent polynomial in $n = r + 1$ variables with*

$$c_{SM}(\mathbf{1}_U) = \sum_{i=0}^r (-1)^i v_i [\mathbb{P}^{r-i}] \in A_*(\mathbb{P}^n).$$

If g is nondegenerate, then

$$v_i = MV_n(\underbrace{\Delta, \dots, \Delta}_{r-i}, \underbrace{\Delta_g, \dots, \Delta_g}_{i+1}), \quad i = 0, \dots, r.$$

In particular, the maximum likelihood degree of U is equal to the normalized volume

$$v_r = (-1)^r \int c_{SM}(\mathbf{1}_U) = \text{Volume}(\Delta_g).$$

Proof. In any case, $c_{SM}(\mathbf{1}_{(\mathbb{C}^*)^n})$ is the fundamental class of the ambient smooth and complete toric variety. Therefore we may assume that U is nonempty and compute the CSM class of $(\mathbb{C}^*)^n \setminus U$ instead.

Fix a sufficiently fine subdivision of the fan of \mathbb{P}^n on which the support function of Δ_g is piecewise linear. We may assume that the corresponding toric variety X is smooth and the closure V of U in X has simple normal crossings with $D := X \setminus (\mathbb{C}^*)^n$ [Hov77, Section 2]. Note in this case that

$$c_{SM}(\mathbf{1}_{(\mathbb{C}^*)^n}) - c_{SM}(\mathbf{1}_U) = c(\Omega_X^1(\log D \cup V)^\vee) \cap [X] \in A_*(X).$$

In order to compute the right-hand side, we use the Poincaré-Leray residue map

$$\text{rés} : \Omega_X^1(\log D \cup V) \longrightarrow \mathcal{O}_V, \quad \eta \cdot d\log(z) + \psi \longmapsto \eta|_V,$$

where z is a local defining equation for V , and ψ is a rational differential one-form which does not have poles along V . The restriction of η on V is uniquely and globally determined, and in particular it does not depend on the choice of z . Note that the residue map fits into the exact sequence

$$0 \longrightarrow \Omega_X^1(\log D) \longrightarrow \Omega_X^1(\log D \cup V) \xrightarrow{\text{rés}} \mathcal{O}_V \longrightarrow 0.$$

Since $\Omega_X^1(\log D)$ is a trivial vector bundle, the Whitney sum formula shows that

$$c(\Omega_X^1(\log D \cup V)^\vee) \cap [X] = \sum_{i=0}^n (-1)^i c_1(\mathcal{O}_X(V))^i \cap [X].$$

Therefore, by the projection formula applied to the birational map $X \rightarrow \mathbb{P}^n$, we have

$$c_{SM}(\mathbf{1}_{(\mathbb{C}^*)^n}) - c_{SM}(\mathbf{1}_U) = \sum_{i=0}^n (-1)^i (H^{n-i} \cdot V^i) [\mathbb{P}^{n-i}] \in A_*(\mathbb{P}^n),$$

where H is the pullback of a hyperplane in \mathbb{P}^n .

It remains to show that the intersection product $(H^{n-i} \cdot V^i)$ is equal to the mixed volume $MV_n(\underbrace{\Delta, \dots, \Delta}_{n-i}, \underbrace{\Delta_g, \dots, \Delta_g}_i)$. For this one computes the divisor of the rational function g on X :

$$\text{Div}(g) = V + \sum_{\rho} \text{ord}_{D_\rho}(g) D_\rho = V + \sum_{\rho} \psi(u_\rho) D_\rho.$$

Here u_ρ is the primitive ray generator of a ray ρ , D_ρ is the torus invariant prime divisor of X corresponding to ρ , and ψ is the support function of Δ_g . It follows that V is linearly equivalent to the torus invariant divisor

$$V_\infty := - \sum_{\rho} \psi(u_\rho) D_\rho.$$

Therefore, by [Ful93, Section 5.4], we have

$$(H^{n-i} \cdot V^i) = (H^{n-i} \cdot V_\infty^i) = \text{MV}_n(\underbrace{\Delta, \dots, \Delta}_{n-i}, \underbrace{\Delta_g, \dots, \Delta_g}_i).$$

□

Remark 4.2. In this case, log-concavity of the v_i discussed in Remark 3.12 is the Alexandrov-Fenchel inequality on the mixed volume of convex bodies [Sch93, Section 6.3].

Remark 4.3. Let \mathcal{V} be a subvariety of the projective space with the homogeneous coordinates p_0, p_1, \dots, p_n . In the statistical setting of [HKS05], one computes the maximum likelihood degree of the very affine variety

$$\mathcal{U} := \{x \in \mathcal{V} \mid p_0 p_1 \cdots p_n (p_0 + p_1 + \cdots + p_n) \neq 0\} \subset \mathbb{P}^n.$$

Suppose \mathcal{V} is the closure of the hypersurface in $(\mathbb{C}^*)^n$ defined by a sufficiently general Laurent polynomial with the given Newton polytope. In the notation used in the proof of Theorem 4.1, we have the residue exact sequence

$$0 \longrightarrow \Omega_X^1(\log D \cup V) \longrightarrow \Omega_X^1(\log D \cup V \cup H) \xrightarrow{\text{rés}} \mathcal{O}_H \longrightarrow 0,$$

where H is the pullback of the hypersurface $\{p_0 + p_1 + \cdots + p_n = 0\}$ in \mathbb{P}^n . Therefore, by the Whitney sum formula,

$$\text{ML}(\mathcal{U}) = v_1 + \cdots + v_n.$$

When the Newton polytope is d -th multiple of the standard simplex Δ , we have

$$\text{ML}(\mathcal{U}) = d + \cdots + d^n = d \cdot \frac{d^n - 1}{d - 1}.$$

This is the formula of [HKS05, Theorem 6].

The remainder of this section is devoted to an application of Theorem 3 to the geometry of projective hypersurfaces. We recall from [Huh12a] the needed properties of CSM classes of projective hypersurfaces. Let h be a nonconstant homogeneous polynomial in $n + 1$ variables

$$h = \sum_{\mathbf{u}} c_{\mathbf{u}} \mathbf{z}^{\mathbf{u}} \in \mathbb{C}[z_0, \dots, z_n].$$

We are interested in the CSM class of the smooth affine variety

$$D(h) = \{h \neq 0\} \subset \mathbb{P}^n.$$

The CSM class of $D(h)$ can be viewed as a projective version of the Milnor number for h , and it is determined by the graph of the gradient map

$$\text{grad}(h) : \mathbb{P}^n \dashrightarrow \mathbb{P}^n, \quad \text{grad}(h) = \left(\frac{\partial h}{\partial z_0} : \cdots : \frac{\partial h}{\partial z_n} \right).$$

The graph $\Gamma_h \subset \mathbb{P}^n \times \mathbb{P}^n$ defines an element in the homology group, which we write as

$$[\Gamma_h] = \sum_{i=0}^n \mu^i(h) [\mathbb{P}^{n-i} \times \mathbb{P}^i] \in A_n(\mathbb{P}^n \times \mathbb{P}^n).$$

For any nonconstant h ,

- $\mu^0(h)$ is one,
- $\mu^1(h)$ is one less than the reduced degree of h ,
- $\mu^n(h)$ is the degree of the gradient map of h , and in general,
- $\mu^i(h)$ is the number of i -dimensional cells in a CW-model of $D(h)$ [Huh12a, Theorem 9].

The CSM class of $D(h)$ is given by the formula

$$c_{SM}(\mathbf{1}_{D(h)}) = \sum_{i=0}^n (-1)^i \mu^i(h) H^i (1 + H)^{n-i} \in A_*(\mathbb{P}^n),$$

where H is the class of a hyperplane in \mathbb{P}^n [Alu03, Theorem 2.1].

Our goal is to show that almost all homogeneous polynomials h with the given Newton polytope Δ_h define a projective hypersurface with the same Milnor numbers $\mu^i(h)$, and there is an explicit formula computing these numbers from Δ_h . This proves an assertion left unjustified in [Huh12a, Example 17].

Let us call a convex polytope Δ_* in \mathbb{R}^{n+1} *homogeneous* if it lies in an affine hyperplane perpendicular to the vector $(1, \dots, 1)$. We normalize the n -dimensional mixed volume for homogeneous polytopes so that the n -dimensional standard simplex Δ_\circ in \mathbb{R}^{n+1} has the unit volume.

For a homogeneous convex polytope Δ_* in \mathbb{R}^{n+1} , set

$$v_i(\Delta_*) := \text{MV}_n(\underbrace{\Delta_\circ, \dots, \Delta_\circ}_{n-i}, \underbrace{\Delta_*, \dots, \Delta_*}_i), \quad i = 0, \dots, n.$$

Note that v_0 vanishes exactly when Δ_* is empty. We extend [Kou76, Définition 1.7] and define the *mixed Newton numbers* of Δ_* by the formula

$$\nu_i(\Delta_*) = V_{n,i} - V_{n-1,i-1} + \dots + (-1)^i V_{n-i,0}, \quad i = 0, \dots, n,$$

where $V_{k,l}$ is the sum of the k -dimensional mixed volumes v_l of the intersections of Δ_* with all possible $(k+1)$ -dimensional coordinate planes.

A homogeneous polynomial is said to be *nondegenerate* if its dehomogenization with respect to a variable is nondegenerate as a Laurent polynomial. Almost all homogeneous polynomials with the given Newton polytope are nondegenerate.

COROLLARY 4.4. *If h is a nondegenerate homogeneous polynomial in $(n+1)$ variables, then*

$$c_{SM}(\mathbf{1}_{D(h)}) = \sum_{i=0}^n (-1)^i \nu_i(\Delta_h) [\mathbb{P}^{n-i}] \in A_*(\mathbb{P}^n).$$

In particular, the degree of the gradient map of h is the sum

$$\mu^n(h) = \sum_{i=0}^n \nu_i(\Delta_h).$$

Proof. Write I for a proper subset of $\{0, 1, \dots, n\}$. We use the decomposition

$$D(h) = \coprod_I D(h) \cap \mathbb{T}_I, \quad \mathbb{T}_I = \{(z_0 : \dots : z_n) \in \mathbb{P}^n \mid z_i = 0, i \in I\}.$$

Let h_I be the homogeneous polynomial obtained from h by setting the variable z_i to zero for $i \in I$. Since every face of the Newton polytope of h_I is a face of the Newton polytope of h , the nondegeneracy of h implies that of h_I . Therefore Theorem 3 computes the CSM class of $D(h) \cap \mathbb{T}_I$ in terms of Δ_{h_I} for every I . Collecting terms of the same dimension in the resulting expressions, we have

$$c_{SM}(\mathbf{1}_{D(h)}) = \sum_I c_{SM}(\mathbf{1}_{D(h) \cap \mathbb{T}_I}) = \sum_{i=0}^n (-1)^i \nu_i(\Delta_h) [\mathbb{P}^{n-i}].$$

□

This shows that many delicate examples discovered in classical projective geometry have a rather simple combinatorial origin. As an example, we consider Hesse's claim which states to the effect that the degree of the gradient map of a homogeneous polynomial is zero if and only if the corresponding projective hypersurface is a cone [Hes51, Hes59]. Hesse's claim is true up to dimension 3, but counterexamples were found by Gordan and Noether in dimension 4 [GN76]. Below is a nondegenerate counterexample with the given degree $d \geq 3$:

$$h = z_3^{d-1} z_0 + z_3^{d-2} z_4 z_1 + z_4^{d-1} z_2.$$

The actual values of the coefficients are irrelevant, so long as they are nonzero; all the polynomials define the same hypersurface up to a linear change of coordinates.

Remark 4.5. Note that we do not have a hypothesis corresponding to *commode* of [Kou76, Définition 1.5]. No positivity property of the Newton numbers should be expected in this generality.

Example 4.6. Homaloidal polynomial is a homogeneous polynomial h whose gradient map is a birational transformation of \mathbb{P}^n . The property of being homaloidal depends only on the set $\{h = 0\} \subseteq \mathbb{P}^n$ [DP03]. We check Corollary 4.4 for three nondegenerate homaloidal plane curves.

1. The nonsingular conic $\{h = x^2 + y^2 + z^2 = 0\} \subset \mathbb{P}^2$: In this case,

$$\begin{aligned} \mu^2(h) &= V_{2,0} + (V_{2,1} - V_{1,0}) + (V_{2,2} - V_{1,1} + V_{0,0}) \\ &= 1 + (2 - 3) + (4 - 6 + 3) \\ &= 1. \end{aligned}$$

2. The union of three nonconcurrent lines $\{h = xyz = 0\} \subset \mathbb{P}^2$: In this case,

$$\begin{aligned} \mu^2(h) &= V_{2,0} + (V_{2,1} - V_{1,0}) + (V_{2,2} - V_{1,1} + V_{0,0}) \\ &= 1 + (0 - 0) + (0 - 0 + 0) \\ &= 1. \end{aligned}$$

3. The union of the conic and its tangent $\{h = zy^2 + x^2y = 0\} \subset \mathbb{P}^2$: In this case,

$$\begin{aligned} \mu^2(h) &= V_{2,0} + (V_{2,1} - V_{1,0}) + (V_{2,2} - V_{1,1} + V_{0,0}) \\ &= 1 + (2 - 2) + (0 - 0 + 0) \\ &= 1. \end{aligned}$$

Dolgachev proved in [Dol00] that any homaloidal plane curve is equal to one of the above, up to a linear change of coordinates of \mathbb{P}^2 . It would be interesting to extend Dolgachev's list and, as a first step, classify nondegenerate homaloidal surfaces in \mathbb{P}^3 .

Example 4.7. The modern study of homaloidal polynomials began with the work of Ein and Shepherd-Barron, and also of Etingof, Kazhdan, and Polishchuk, who observed that the relative invariant of a regular prehomogeneous vector space is homaloidal [ES89, EKP02]. The degree of these homaloidal polynomials are bounded in terms of the ambient dimension. In contrast to the case of relative invariants and of plane curves, Ciliberto, Russo, and Simis showed that there are irreducible homaloidal hypersurfaces of the given degree $d \geq 2n - 3$ and the ambient dimension $n \geq 3$ [CRS08, Theorem 3.13]. Their construction relies on an ingenious argument concerning projective duals of rational scroll surfaces.

Corollary 4.4 provides a cheap way of producing projective hypersurfaces with interesting numerical properties (perhaps with the aid of a computer), and in particular homaloidal polynomials. Once the correct form of the equation is guessed, it is often not hard to check by hand that the example has the desired property. We demonstrate this by showing that there are irreducible homaloidal hypersurfaces of the given degree $d \geq 3$ and the ambient dimension $n \geq 3$.

Among many explicit examples of this type, we present a particularly transparent one:

$$h = z_0 z_n^{d-1} + z_1 z_2 z_n^{d-2} + z_2^d + (z_3^2 + z_4^2 + \cdots + z_{n-1}^2) z_n^{d-2}.$$

The actual values of the coefficients are irrelevant, so long as they are nonzero; all the polynomials define the same hypersurface up to a linear change of coordinates. By the Bertini theorem, h is nondegenerate and irreducible. The system of equations

$$y_0 = \frac{\partial h}{\partial z_0}, \dots, y_n = \frac{\partial h}{\partial z_n}$$

is triangular (of De Jonquières type), defining an isomorphism between the affine spaces $z_n \neq 0$ and $y_0 \neq 0$. Therefore h is homaloidal for any $d \geq 3$ and $n \geq 3$. We note that Fassarella and Medeiros constructed (necessarily reducible) homaloidal hypersurfaces with the given degree $d \geq 3$ and the ambient dimension $n \geq 3$, using an idea based on the CSM class [FM12, Example 5.1].

REFERENCES

- Alu99 Paolo Aluffi, *Differential forms with logarithmic poles and Chern-Schwartz-MacPherson classes of singular varieties*, Comptes Rendus de l'Académie des Sciences. Série I. Mathématique **329** (1999), 619–624.
- Alu03 Paolo Aluffi, *Computing characteristic classes of projective schemes*, Journal of Symbolic Computation **35** (2003), 3–19.
- Alu05 Paolo Aluffi, *Characteristic classes of singular varieties*, Topics in Cohomological Studies of Algebraic Varieties, 1–32, Trends in Mathematics, Birkhäuser, Basel, 2005.
- Alu06a Paolo Aluffi, *Classes de Chern des variétés singulières, revisitées*, Comptes Rendus Mathématique. Académie des Sciences. Paris **342** (2006), 405–410.
- Alu06b Paolo Aluffi, *Limits of Chow groups, and a new construction of Chern-Schwartz-MacPherson classes*, Pure and Applied Mathematics Quarterly **2** (2006), 915–941.
- Alu12 Paolo Aluffi, *Grothendieck classes and Chern classes of hyperplane arrangements*, International Mathematical Research Notices, to appear.
- CHKS06 Fabrizio Catanese, Serkan Hosten, Amit Khetan, and Bernd Sturmfels, *The maximum likelihood degree*, American Journal of Mathematics **128** (2006), 671–697.
- CRS08 Ciro Ciliberto, Francesco Russo, and Aron Simis, *Homaloidal hypersurfaces and hypersurfaces with vanishing Hessian*, Advances in Mathematics **218** (2008), 1759–1805.

- CDFV11 Daniel Cohen, Graham Denham, Michael Falk, and Alexander Varchenko, *Critical points and resonance of hyperplane arrangements*, Canadian Journal of Mathematics **63** (2011), 1038–1057.
- CLO98 David Cox, John Little, and Donal O’Shea, *Using Algebraic Geometry*, Graduate Texts in Mathematics **185**, Springer, New York, 1998.
- Dam99 James Damon, *Critical points of affine multiforms on the complements of arrangements*, Singularity Theory, 25–33, London Mathematical Society Lecture Note Series **263**, Cambridge University Press, 1999.
- Dam00 James Damon, *On the number of bounding cycles for nonlinear arrangements*, Arrangements - Tokyo 1998, 51–72, Advanced Studies in Pure Mathematics **27**, Kinokuniya, Tokyo, 2000.
- Daw84 Jeremy Dawson, *A collection of sets related to the Tutte polynomial of a matroid*, Graph theory, Singapore 1983, Lecture Notes in Mathematics **1073**, 193–204, Springer, Berlin, 1984.
- DP95 Corrado De Concini and Claudio Procesi, *Wonderful models of subspace arrangements*, Selecta Mathematica. New Series **1** (1995), 459–494.
- Del70 Pierre Deligne, *Équations différentielles à points singuliers réguliers*, Lecture Notes in Mathematics **163**, Springer-Verlag, Berlin-New York, 1970.
- DGS12 Graham Denham, Mehdi Garrousian, and Mathias Schulze, *A geometric deletion-restriction formula*, Advances in Mathematics **230** (2012), 1979–1994.
- DP03 Alexandru Dimca and Stefan Papadima, *Hypersurface complements, Milnor fibers and higher homotopy groups of arrangements*, Annals of Mathematics (2) **158** (2003), 473–507.
- Dol00 Igor Dolgachev, *Polar Cremona transformations*, Michigan Mathematical Journal **48** (2000), 191–202.
- ES89 Lawrence Ein and Nicholas Shepherd-Barron, *Some special Cremona transformations*, American Journal of Mathematics **111** (1989), 783–800.
- EKL06 Manfred Einsiedler, Mikhail Kapranov, and Douglas Lind, *Non-archimedean amoebas and tropical varieties*, Journal für die reine und angewandte Mathematik **601** (2006), 139–157.
- EKP02 Pavel Etingof, David Kazhdan, and Alexander Polishchuk, *When is the Fourier transform of an elementary function elementary?*, Selecta Mathematica. New Series **8** (2002), 27–66.
- FM12 Thiago Fassarella and Nivaldo Medeiros, *On the polar degree of projective hypersurfaces*, Journal of the London Mathematical Society, to appear.
- Ful93 William Fulton, *Introduction to Toric Varieties*, Annals of Mathematics Studies **131**, Princeton University Press, Princeton, NJ, 1993.
- Ful98 William Fulton, *Intersection Theory*, Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics **2**, Springer-Verlag, Berlin, 1998.
- GN76 Paul Gordan and Max Noether, *Ueber die algebraischen Formen, deren Hesse’sche Determinante identisch verschwindet*, Mathematische Annalen **10** (1876), 547–568.
- GP02 Mark Goresky and William Pardon, *Chern classes of automorphic vector bundles*, Inventiones Mathematicae **147** (2002), 561–612.
- Hac08 Paul Hacking, *Homology of tropical varieties*, Collectanea Mathematica **59** (2008), 263–273.
- Hes51 Otto Hesse, *Über die Bedingung, unter welcher eine homogene ganze Function von n unabhängigen Variablen durch lineäre Substitutionen von n andern unabhängigen Variablen auf eine homogene Function sich zurück-führen läßt, die eine Variable weniger enthält*, Journal für die reine und angewandte Mathematik **42** (1851), 117–124.
- Hes59 Otto Hesse, *Zur Theorie der ganzen homogenen Functionen*, Journal für die reine und angewandte Mathematik **56** (1859), 263–269.
- Hog74 Stuart Hoggar, *Chromatic polynomials and logarithmic concavity*, Journal of Combinatorial Theory Series B **16** (1974), 248–254.

- HKS05 Serkan Hoşten, Amit Khetan, and Bernd Sturmfels, *Solving likelihood equations*, Foundations of Computational Mathematics **5** (2005), 389–407.
- Hov77 Askold Hovanskii, *Newton polyhedra and toroidal varieties*, Akademija Nauk SSSR. Funkcional’nyi Analiz i ego Prilozhenija **11** (1977), 56–64.
- Huh12a June Huh, *Milnor numbers of projective hypersurfaces and the chromatic polynomial of graphs*, Journal of the American Mathematical Society **25** (2012), 907–927.
- Huh12b June Huh, *h-vectors of matroids and logarithmic concavity*, preprint, [arXiv:1201.2915](https://arxiv.org/abs/1201.2915).
- Jou83 Jean-Pierre Jouanolou, *Théorèmes de Bertini et Applications*, Progress in Mathematics **42**, Birkhäuser Boston, Boston, MA, 1983.
- Kaw78 Yujiro Kawamata, *On deformations of compactifiable complex manifolds*, Mathematische Annalen **235** (1978), 247–265.
- Kou76 A. G. Kouchnirenko, *Polyèdres de Newton et nombres de Milnor*, Inventiones Mathematicae **32** (1976), 1–31.
- Kwi92 Michał Kwieciński, *Formule du produit pour les classes caractéristiques de Chern-Schwartz-MacPherson et homologie d’intersection*, Comptes Rendus de l’Académie des Sciences. Série I. Mathématique **314** (1992) 625–628,
- Len12 Matthias Lenz, *The f-vector of a realizable matroid complex is strictly log-concave*, Combinatorics, Probability, and Computing, to appear.
- Mac74 Robert MacPherson, *Chern classes for singular algebraic varieties*, Annals of Mathematics (2) **100** (1974), 423–432.
- Nor78 Yoshiki Norimatsu, *Kodaira vanishing theorem and Chern classes for ∂ -manifolds*, Proceedings of the Japan Academy. Series A **54** (1978), 107–108.
- OT92 Peter Orlik and Hiroaki Terao, *Arrangements of Hyperplanes*, Grundlehren der Mathematischen Wissenschaften **300**, Springer-Verlag, Berlin, 1992.
- OT95 Peter Orlik and Hiroaki Terao, *The number of critical points of a product of powers of linear functions*, Inventiones Mathematicae **120** (1995), 1–14.
- PS05 Lior Pachter and Bernd Sturmfels, *Algebraic Statistics for Computational Biology*, Cambridge University Press, New York, 2005.
- Rea68 Ronald Read, *An introduction to chromatic polynomials*, Journal of Combinatorial Theory **4** (1968), 52–71.
- Sai80 Kyoji Saito, *Theory of logarithmic differential forms and logarithmic vector fields*, Journal of the Faculty of Science. University of Tokyo. Section IA. Mathematics **27** (1980), 265–291.
- Sam66 Pierre Samuel, *À propos du théorème des unités*, Bulletin des Sciences Mathématiques (2) **90** (1966), 89–96.
- Sch93 Rolf Schneider, *Convex Bodies: The Brunn-Minkowski Theory*, Encyclopedia of Mathematics and its Applications **44**, Cambridge University Press, Cambridge, 1993.
- Sil96 Roberto Silvotti, *On a conjecture of Varchenko*, Inventiones Mathematicae **126** (1996), 235–248.
- Spe05 David Speyer, *Tropical Geometry*, Ph. D. Thesis, University of California, Berkeley, 2005.
- SS04 David Speyer and Bernd Sturmfels, *The tropical Grassmannian*, Advances in Geometry **4** (2004), 389–411.
- Tev07 Jenia Tevelev, *Compactifications of subvarieties of tori*, American Journal of Mathematics **129** (2007), 1087–1104.
- Var76 Alexander Varchenko, *Zeta-function of monodromy and Newton’s diagram*, Inventiones Mathematicae **37** (1976), 253–262.
- Var95 Alexander Varchenko, *Critical points of the product of powers of linear functions and families of bases of singular vectors*, Compositio Mathematica **97** (1995), 385–401.
- Wel69 Dominic Welsh, *Combinatorial problems in matroid theory*, Combinatorial Mathematics and its Applications (Proc. Conf., Oxford, 1969), 291–306, Academic Press, London, 1971.

THE MAXIMUM LIKELIHOOD DEGREE OF A VERY AFFINE VARIETY

June Huh juneuh@umich.edu

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA